



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bo

Search PubMed

for

Limits

Preview/Index

History

Clipboard

Details

Go Clear

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

1: J Invest Dermatol. 1992 Sep;99(3):283-8.

Related Articles, Links

**Short-term retinoic acid treatment increases in vivo, but decreases in vitro, epidermal transglutaminase-K enzyme activity and immunoreactivity.****Griffiths CE, Rosenthal DS, Reddy AP, Elder JT, Astrom A, Leach K, Wang TS, Finkel LJ, Yuspa SH, Voorhees JJ, et al.**

Department of Dermatology, University of Michigan Medical Center, Ann Arbor 48109-0314.

Epidermal transglutaminase-K is believed to catalyze the covalent linking of loricrin and involucrin to form cross-linked (CE) envelopes. In normal skin, transglutaminase-K is expressed as a band immediately below the stratum corneum, whereas in psoriasis and healing skin its expression is considerably expanded throughout the suprabasal layers. We have investigated whether the hyperproliferative state induced by short-term application of topical retinoic acid is similarly characterized by an increase in transglutaminase-K enzyme activity and immunoreactivity. Retinoic acid (0.1% cream) or vehicle were applied to human skin and occluded for 4 d. Skin biopsies were obtained for measurement of transglutaminase-K and transglutaminase-C activity and immunoreactivity. For comparison, cultured normal human keratinocytes were incubated for 4 d in the presence of 1 microM retinoic acid and the subsequent transglutaminase-K activity and immunoreactivity measured. Transglutaminase-K activity was increased 2.8 times in retinoic acid compared to vehicle-treated skin ( $p$  less than 0.005,  $n = 12$ ) whereas there was no significant difference in transglutaminase-C activity. However, transglutaminase-K mRNA levels were not significantly different between retinoic acid- and vehicle-treated skin. In vehicle-treated skin, transglutaminase-K immunoreactivity was limited to a narrow, substratum corneal band, but was considerably expanded in a diffuse suprabasal pattern in retinoic acid-treated epidermis. In contrast, transglutaminase-K immunostaining was decreased and its enzymatic activity reduced sixfold in retinoic acid-treated keratinocytes ( $p$  less than 0.01,  $n = 4$ ). These results demonstrate that retinoic acid treatment in vivo, in contrast to in vitro, leads to not only increased transglutaminase-K protein expression but also increased enzymatic activity in the absence of detectable increases in mRNA levels. These data, taken with the previously reported lack of in vivo modulation of the differentiation markers keratins 1 and 10 by retinoic acid, indicate that certain aspects of keratinocyte terminal differentiation that are altered in vitro by retinoic acid do not occur in vivo in human skin.

**Publication Types:**

- Clinical Trial
- Randomized Controlled Trial

PMID: 1355099 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text